Journal of Surgical Oncology

Table of Contents

Volume 80, Number 1, May 2002

Guest Editorial: Palliative Surgery: A Modern Perspective
Lawrence D. Wagman 1

Relationship of Clinical and Pathologic Response to Neoadjuvant Chemotherapy and Outcome of Locally Advanced Breast Cancer
Csaba Gajdos, Paul Ian Tartter, Alison Estabrook, Michael A. Gistrak, Shabnam Jaffer, and Ira J. Bleiweiss 4
This study of patients treated with neoadjuvant chemotherapy for locally advanced breast cancers found little evidence that measurable clinical or pathologic changes attributable to chemotherapy predicted survival. Axillary lymph node status, associated with young age, was the most important prognostic indicator in these patients.

Impact of Patient Age on the Outcome of Primary Breast Carcinoma
Byeong-Woo Park, Seung-Il Kim, Eun Kyong Kim, Woo-Ick Yang, and Kyong Sik Lee 12
This study showed that a patient aged younger than 35 is an independent prognostic factor. We believe there may be a high-risk subset of breast cancers with respect to recurrence and that a higher proportion of breast cancers arising in women younger than 35 years old fall into this high-risk subset, which requires aggressive treatments based on the tumor biology.

Primary Non-Hodgkin Lymphoma of the Breast: The Mayo Clinic Experience
William W. Wong, Steven E. Schild, Michele Y. Halyard, and Paula J. Schomberg 19
This study provides a summary of the natural history, treatment results, and prognostic factors for patients with primary breast non-Hodgkin lymphoma. Recommendations for management of these patients are suggested.

Commentary
Cary S. Kaufman 26

Ex Vivo Sentinel Lymph Node Biopsy in Colorectal Cancer: A Feasibility Study
T.L. Fitzgerald, M.A. Khalifa, M. Al Zahrani, C.H.L. Law, and A.J. Smith 27
Sentinal lymph node (SLN) biopsy may improve staging of colorectal cancer. We tested the feasibility of ex vivo SLN dissection.

Commentary
Anton J. Bilchik 33

Microsatellite Analysis of Fecal DNA for Colorectal Cancer Detection
Minori Koshiji, Yasuhiro Yonekura, Takamichi Saito, and Kazuhiko Yoshioka 34
In an effort to develop a noninvasive detection test for colorectal cancer, we performed microsatellite analyses using blood, surgical specimens, oral rinse, and stool samples.

Formalin Instillation for Refractory Radiation-Induced Hemorrhagic Proctitis
Pedro Luna-Pérez and Saúl E. Rodríguez-Ramírez 41
Radiation proctitis is a common complication after pelvic irradiation. One to five percent of these patients will develop intractable or massive hemorrhagic radiation proctitis. Severe hemorrhagic radiation proctitis is a tremendous therapeutic challenge. Twenty patients who failed with administration of topical steroids and/or mesalazine were treated with 500 mL of 4% formalin instilled into the rectum in 50-mL aliquots with an overall success rate of 90% median follow-up of 20 months.

Significance of Postoperative Serum Thyroglobulin Levels in Patients with Papillary and Follicular Thyroid Carcinomas
Jen-Der Lin, Miau-Ju Huang, Brend Ray-Sea Hsu, Tzu-Chieh Chao, Chuen Hsueh, Feng-Hsuan Liu, Miaw-Jene Liou, and Hsiao-Fen Weng 45
Thyroglobulin is one of the important differentiation markers in thyroid follicular epithelial cells. Postoperation serum levels of thyroglobulin can be used as a prognostic indicator in patients with papillary and follicular thyroid cancer.

Mutations in the von Hippel-Lindau (VHL) Gene Refine Differential Diagnostic Criteria in Renal Cell Carcinoma
Nandita Barnabas, Mitual B. Amin, Kirit Pindolia, Reshma Nanavati, Mahul B. Amin, and Maria J. Worsham 52
A cohort of paraffin embedded sporadic Renal Cell Carcinoma (RCC) specimens revealed VHL mutations in the elongin binding and 5'UTR regions. Mutations in chromophobe RCC are a novel finding.

How I Do It: Costal Chondritis After Thoracoabdominal Esophagectomy: How To Prevent It
J. Edward M. Young, John D. Miller, and John D. Urschel 61

Commentary
Constantine P. Karakousis 63

Letter To The Editor: Hyperthermic Isolated Hepatic Perfusion for the Treatment of Unresectable Cancers Confined to the Liver
Nancy M. Carroll and H. Richard Alexander 64

Reply
Paul Moroz, Stephen K. Jones, and Bruce N. Gray 66

Volume 80, Number 2, June 2002

Guest Editorial: Palliative Liver Resections
Kenneth K. Tanabe 69

Concentrated Preoperative Radiotherapy for Resectable Gastric Cancer: 20-Years Follow-Up of a Randomized Trial
Vitali Skoropad, Boris Berdov, and Vasili Zagrebin 72
Concentrated preoperative radiotherapy at a dose of 20 Gy did not significantly increase survival of patient with gastric cancer. However, a trend toward improved results was seen that supports further investigations of adjuvant radiation therapy in gastric cancer.

Expression of Cyclooxygenase-2 in Primary and Remnant Gastric Carcinoma: Comparing It With p53 Accumulation, *Helicobacter pylori* Infection, and Vascular Endothelial Growth Factor Expression
Atsushi Kawabe, Yutaka Shimada, Shigeki Uchida, Masato Maeda, Seiji Yamasaki, Masayuki Kato, Yosuke Hashimoto, Gakuji Ohshio, Masao Matsumoto, and Masayuki Imamura 79
In a series of 91 gastric carcinoma cases, COX-2 (Cyclooxygenase-2) was frequently expressed and there was significant correlation between COX-2 and p53 accumulation. The positive rate of COX-2 was significantly higher in noncancerous adjacent mucosa in remnant gastric cancer cases than in primary gastric cancer cases.

Metastatic Patterns of Extremity Myxoid Liposarcoma and Their Outcome
Susanne H. Estourgie, Gunnlaugur P. Nielsen, and Mark J. Ott 89
Extremity myxoid liposarcomas have an unusually high predilection for intra-abdominal, retroperitoneal, and paraspinal metastases. After treatment of the primary tumor, these patients should be followed with periodic chest X-ray and abdominal/pelvic computed tomography scans and consideration should be given to surgical and adjuvant treatment of metastatic disease when appropriate.

Commentary: Metastatic Patterns of Extremity Liposarcoma and Their Outcome
Peter W. T. Pisters and Vernon K. Sondak 94

Desmoid Tumors: A Novel Approach for Local Control
Christopher R. Baliski, Walley J. Temple, Keith Arthur, and Norman S. Schachar 96
The results of a prospective cohort study using neoadjuvant chemoradiotherapy in the treatment of primary and recurrent desmoid tumors was analyzed. Preoperative concomitant low dose doxyrubicin (90 mg over 3 days) and radiation (300cGy × 10 days), followed by surgery was completed in 13 resectable patients. With a median follow up of 71 months (range 22–109), there were 2 local recurrences (15% local recurrence). One patient developed lesions in other anatomic sites from multifocal disease, which was not considered recurrent disease.

Multivisceral Resection for Colon Cancer
Pedro Luna-Pérez, Saúl E. Rodríguez-Ramírez, Marcos Gutiérrez de la Barrera, Moisés Zeferino, and Sonia Labastida 100
Colon cancer with neoplasic infiltration to neighboring organs occurs in approximately 5–10% of patients with colon cancer. En bloc surgical resection is the standard of care for this group of patients. In the current analysis, risk factors for cancer-related failure are related with neoplasic infiltration to neighboring organs (T4), age > 50 years, and lymph node metastases. In T4 colon cancer, the recurrence pattern was found at local, peritoneal, and distant sites.

Comparison of Matrix Metalloproteinase Expression Between Primary Tumors With or Without Liver Metastasis in Pancreatic and Colorectal Carcinomas
Yoshito Matsuyama, Sonshin Takao, and Takashi Aikou 105
Matrix metalloproteinases (MMPs) are known to play an important role in carcinoma cell invasion and hematogenous metastasis by mediating the degradation of the extracellular matrix. We investigated the relationship between MMP-2 and -9 enzymatic activities and liver metastases in human pancreatic and colon carcinomas using xenograft tumors in nude mice.

Large Bowel Lymphoma: An Analysis of Prognostic Factors and Therapy in 53 Patients
Agustin Avilés, Natividad Neri, and Judith Huerta-Guzmán 111

An retrospective analysis of 53 patients with large bowel lymphoma shown that surgery followed by adjuvant chemotherapy is the treatment of choice and that prognostic factors can not be identified.

How I Do It: Exteriorized Small Bowel as an Abdomino-Pelvic Partition
Paul H. Sugarbaker 116

Volume 80, Number 3, July 2002

Guest Editorial: The TNM System: Our Language for Cancer Care
Frederick L. Greene and Leslie H. Sobin 119

Sole Brachytherapy of the Tumor Bed After Conservative Surgery for T1 Breast Cancer: Five-Year Results of a Phase I–II Study and Initial Findings of a Randomized Phase III Trial
Csaba Polgár, Zoltán Sulyok, János Fodor, Zsolt Orosz, Tibor Major, Zoltán Takácsi-Nagy, László C. Mangel, András Somogyi, Miklós Kásler, and György Németh 121
Experience from prospective clinical trials with sole tumor bed radiotherapy after breast-conserving surgery was reported, including the 5-year results of a phase I–II study and initial findings of a phase III trial comparing the efficacy of tumor bed irradiation alone with conventional whole breast radiotherapy. A 4-day course of sole brachytherapy with careful patient selection and adequate quality assurance appears to be a feasible alternative to whole breast radiotherapy.

Commentary
Robert J. Dowsett and Scott H. Kurtzman 129

Sentinel Node Biopsy for Breast Cancer: Does the Number of Sentinel Nodes Removed Have an Impact on the Accuracy of Finding a Positive Node?
Peter Schrenk, Wolfgang Rehberger, Andreas Shamiyeh, and Wolfgang Wayand 130
The number of sentinel lymph nodes removed during sentinel node biopsy was not related to patient and tumor factors or to parameters related to the technique of sentinel node biopsy. The pathologic status of the axilla was independently determined by removal of the first or first and second sentinel node in 99% of patients, and the removal of more than three sentinel nodes did not increase the accuracy of finding a positive sentinel node.

Influence of Environment and Healthcare Structures on the Survival of Patients With Colorectal Cancer: A French Population-Based Study
Frédéric Borie, Jean-Pierre Daurs, Bertrand Millat, Pierre Puch, and Brigitte Tretarre 137
Heterogeneity and inequality persist in the distribution of, the treatment for, and the survival of patients with colorectal cancer based on the type of healthcare facility and the living area.

Hepatic Arterial Infusion of 5-Fluorouracil and Cisplatin for Unresectable or Recurrent Hepatocellular Carcinoma With Tumor Thrombus of the Portal Vein
Toshiyuki Itamoto, Hideki Nakahara, Hirotaka Tashiro, Naoki Haruta, Toshimasa Asahara, Akira Naito, and Katsuhide Ito 143
The study was designed to evaluate the efficacy of hepatic arterial infusion of 5-fluorouracil (5-FU) and cisplatin (CDDP) for unresectable or recurrent hepatocellular carcinoma (HCC) with tumor thrombus of the trunk or first branches of the portal vein (PVTT).

Tumor Response to Arterial Embolization Hyperthermia and Direct Injection Hyperthermia in a Rabbit Liver Tumor Model
Paul Moroz, Stephen K. Jones, and Bruce N. Gray 149
The objective of this study was to compare the response of liver tumors to arterial embolization hyperthermia (AEH) and direct injection hyperthermia (DIH).

Review Article: Diagnosis and Management of Patients With Thyroid Nodules
Walter Lawrence, Jr. and Brian J. Kaplan 157

How I Do It: Surgical Lesioning of Splanchnic Nerves Using Wet Needle Radiofrequency Thermoablation
Serge Evrard 171

How I Do It: Radio-Guided Parathyroidectomy for Primary Hyperparathyroidism Combined With Video-Assisted Surgery Using the Solid-State, Multi-Crystal Gamma Camera
Wataru Kitagawa, Kazuo Shimizu, Shin-Ichiro Kumita, Haruki Akasu, Tatsuo Kumazaki, and Shigeo Tanaka 173

Letter to the Editor: Treatment of Neoplastic Diseases of the Sacrum
Markus Wenger and Thomas Marc Markwalder 176

Volume 80, Number 4, August 2002

Guest Editorial: Surgical Quality in Trials of Adjuvant Cancer Therapy
Scott A. Hundahl 177

Outcomes after Emergency Surgery for Gastric Perforation or Severe Bleeding in Patients with Gastric Cancer
Yuichi Kasakura, Jaffer A. Ajani, Fumiro Mochizuki, Yukie Morishita, Masashi Fujii, and Tadatoshi Takayama 181
This study was undertaken to clarify the clinicopathologic characteristics of gastric-cancer patients with free perforation and major bleeding and to determine optimum management of these conditions.

Predictors of Nodal Metastasis in Salivary Gland Cancer
Hakan Korkmaz, George H. Yoo, Wei Du, Eitan Hocwald, Jose E. Otero-Garcia, N. Volkan Adsay, Terry Shibuya, and John R. Jacobs 186
The results of a study of 40 patients suggest that clinical neck examination is a reliable predictor of regional metastasis in patients with major salivary gland cancer.

Analysis of Gene Expression in Egr-1 Transfected Human Fibrosarcoma Cells
Ya-Guang Liu, Jacob Gano, Ruochun Huang, Ying Lin, Shen-Ming Wang, and Ruo-Pan Huang 190
Because Egr-1 is a transcription factor, tumor suppressor function is likely to be involved in its transcriptional activity. We therefore analyzed Egr-1-regulated genes by means of cDNA microarray.

Establishment and Characterization of OCUT-1, an Undifferentiated Thyroid Cancer Cell Line Expressing High Level of Telomerase
Kana Ogisawa, Naoyoshi Onoda, Tetsuro Ishikawa, Chiemi Takenaka, Masaaki Inaba, Yoshinari Ogawa, and Kosei Hirakawa-YS Chung 197
Establishment and characterization of an undifferentiated thyroid cancer cell line, OCUT-1, derived from the surgical specimen of a case with advanced undifferentiated thyroid cancer are reported.

Review Article: Genetic Profiling of Colon Cancer
Holly L. Neibergs, David W. Hein, and John S. Spratt 204

How I Do It: Radical Resection of a Malignant Mesenchymoma With Hypogastric Artery Transposition
Andrew C. Bourne, Larry W. Kraiss, Joseph A. Holden, Mary C. Mone, and Richard G. Barton 214

How I Do It: Minimally Invasive Video-Assisted Thyroidectomy and Lymphadenectomy for Micropapillary Carcinoma of the Thyroid
Yoshifumi Ikeda, Hiroshi Takami, Yuzo Sasaki, Junichi Takayama, Shigenao Kan, and Masanori Niimi 218

How I Do It: Computer Graphic to Design the Approach for Resection of Esthesioneuroblastoma
Susan Burekhovich and Bimal C. Ghosh 222

Letter to the Editor: Reoperation Combined With Intraoperative Hyperthermic Intrathoracic Perfusion Chemotherapy for Pleural Recurrence of Thymoma
Eelco de Bree, Serge van Ruth, Emiel J. Th. Rutgers, and Frans A. N. Zoetmulder 224

Author Index to Volume 80 226

Subject Index to Volume 80 228